

# Quotient Category and Sub-Category

Start with homotopy category

Hao Fan

Undergraduate  
Mathematics and Applied mathematics, T-2001

Course Report, March 2023



# Table of Contents

1 Quotient Category

2 Sub-Category



# Recall

We have studied a special category ( $\text{hTop}$ ), defined as follows:

$$\text{Ob}(\text{hTop}) := \text{Ob}(\text{Top}),$$

$$\text{Hom}_{(\text{hTop})}(X, Y) := \text{Hom}_{(\text{Top})}(X, Y)/\simeq,$$

where  $\simeq$  satisfies:

$$f_0 \simeq f_1, g_0 \simeq g_1 \implies g_0 \circ f_0 \simeq g_1 \circ f_1.$$



# Table of Contents

1 Quotient Category

2 Sub-Category



# Quotient Category

As a generalization of (hTop):

## Definition

Let a category  $\mathcal{C}$  and (a family of) equivalence relations  $\stackrel{A,B}{\sim}$  on each set  $\text{Hom}_{\mathcal{C}}(A, B)$ , where  $A, B \in \text{Ob}(\mathcal{C})$  be given. We define the **quotient category** w.r.t  $\sim$ , denoted by  $\mathcal{C}/\sim$ , as follows:

$$\text{Ob}(\mathcal{C}/\sim) := \text{Ob}(\mathcal{C}),$$

$$\text{Hom}_{\mathcal{C}/\sim}(A, B) := \text{Hom}_{\mathcal{C}}(A, B)/\stackrel{A,B}{\sim},$$

where  $\simeq$  satisfies:  $f_0 \simeq f_1, g_0 \simeq g_1 \implies g_0 \circ f_0 \simeq g_1 \circ f_1$ . It has the identity automatically and the associativity holds.



# Quotient Category

## Remark

There is a “natural” full functor:  $Q: \mathcal{C} \rightarrow \mathcal{C}/\sim$ .



# Quotient Category

## Remark

There is a “natural” full functor:  $Q: \mathcal{C} \rightarrow \mathcal{C}/\sim$ .

## Corollary

*If  $A \cong B$  in the category  $\mathcal{C}$ , then  $A \cong B$  in  $\mathcal{C}/\sim$ .*



# Examples of Quotient Categories

## Example

The category  $(hTop) = (Top)/\simeq$ , where  $\simeq$  is the homotopy relation.

## Example

The category  $(\text{Lin}_{\mathbb{C}})$ , with equivalence relations  $\sim$  defined as follows: for  $f, g \in \mathcal{L}(X, Y)$ ,  $f \sim g \iff \exists k \in \mathbb{C} \setminus \{0\}$  such that  $f = kg$ . Then we have a quotient category  $(\text{Lin}_{\mathbb{C}})/\sim$ .



# Table of Contents

1 Quotient Category

2 Sub-Category



# Sub-Category

## Definition

Let a category  $\mathcal{C}$  be given. If another category  $\mathcal{C}'$  satisfies:

- ①  $\text{Ob}(\mathcal{C}') \subseteq \text{Ob}(\mathcal{C})$ ;
- ②  $\forall A, B \in \text{Ob}(\mathcal{C}') : \mathcal{C}'(A, B) := \text{Hom}_{\mathcal{C}'}(A, B) \subseteq \mathcal{C}(A, B)$ ;

# Sub-Category

## Definition

Let a category  $\mathcal{C}$  be given. If another category  $\mathcal{C}'$  satisfies:

- ①  $\text{Ob}(\mathcal{C}') \subseteq \text{Ob}(\mathcal{C})$ ;
- ②  $\forall A, B \in \text{Ob}(\mathcal{C}') : \mathcal{C}'(A, B) := \text{Hom}_{\mathcal{C}'}(A, B) \subseteq \mathcal{C}(A, B)$ ;
- ③ compositions in  $\mathcal{C}'$  are birestrictions of compositions in  $\mathcal{C}$ , i.e. the following diagram commutes;

$$\begin{array}{ccc} \mathcal{C}'(A, B) \times \mathcal{C}'(B, C) & \longrightarrow & \mathcal{C}'(A, C) \\ \downarrow & & \downarrow \\ \mathcal{C}(A, B) \times \mathcal{C}(B, C) & \longrightarrow & \mathcal{C}(A, C) \end{array}$$

- ④  $\forall A \in \text{Ob}(\mathcal{C}')$ , the identity map of  $A$  (in  $\mathcal{C}$ ), i.e.  $1_A^{\mathcal{C}}$  lies in  $\mathcal{C}'(A, A)$ ;
- then we say  $\mathcal{C}'$  is a **sub-category** of  $\mathcal{C}$ , denoted by  $\mathcal{C}' \subseteq \mathcal{C}$ .

# Sub-Category

## Remark

To describe a sub-category of  $\mathcal{C}$ , it suffices to tell its object class and morphism sets. Since all the others are determined by  $\mathcal{C}$ .



# Sub-Category

## Remark

To describe a sub-category of  $\mathcal{C}$ , it suffices to tell its object class and morphism sets. Since all the others are determined by  $\mathcal{C}$ .

## Remark

There is a “natural” faithful functor  $I: \mathcal{C}' \rightarrow \mathcal{C}$ .



# Sub-Category

## Remark

To describe a sub-category of  $\mathcal{C}$ , it suffices to tell its object class and morphism sets. Since all the others are determined by  $\mathcal{C}$ .

## Remark

There is a “natural” faithful functor  $I: \mathcal{C}' \rightarrow \mathcal{C}$ .

## Corollary

*If  $A \cong B$  in the category  $\mathcal{C}'$ , then  $A \cong B$  in  $\mathcal{C}$ .*



# Sub-Category

## Remark

To describe a sub-category of  $\mathcal{C}$ , it suffices to tell its object class and morphism sets. Since all the others are determined by  $\mathcal{C}$ .

## Remark

There is a “natural” faithful functor  $I: \mathcal{C}' \rightarrow \mathcal{C}$ .

## Corollary

*If  $A \cong B$  in the category  $\mathcal{C}'$ , then  $A \cong B$  in  $\mathcal{C}$ .*

## Definition

A sub-category is said to be **full**, if the functor  $I: \mathcal{C}' \rightarrow \mathcal{C}$  is full.



# Sub-Category

The following are sub-categories relevant to our familiar categories:



# Sub-Category

The following are sub-categories relevant to our familiar categories:

## Examples

- ①  $(\text{FinSet}) \subseteq (\text{Set})$ , it is full;



# Sub-Category

The following are sub-categories relevant to our familiar categories:

## Examples

- ①  $(\text{FinSet}) \subseteq (\text{Set})$ , it is full;
- ②  $(\text{FinLin}_{\mathbb{C}}) \subseteq (\text{Lin}_{\mathbb{C}})$ , it is full;



# Sub-Category

The following are sub-categories relevant to our familiar categories:

## Examples

- ①  $(\text{FinSet}) \subseteq (\text{Set})$ , it is full;
- ②  $(\text{FinLin}_{\mathbb{C}}) \subseteq (\text{Lin}_{\mathbb{C}})$ , it is full;
- ③  $(\text{Ab}) \subseteq (\text{Grp})$ , it is full;



# Sub-Category

The following are sub-categories relevant to our familiar categories:

## Examples

- ①  $(\text{FinSet}) \subseteq (\text{Set})$ , it is full;
- ②  $(\text{FinLin}_{\mathbb{C}}) \subseteq (\text{Lin}_{\mathbb{C}})$ , it is full;
- ③  $(\text{Ab}) \subseteq (\text{Grp})$ , it is full;
- ④  $(\text{Haus}) \subseteq (\text{Top})$ , it is full;



# Sub-Category

## Example

Let a fixed category  $\mathcal{C}$  satisfying  $\text{Aut}(X_0, Y_0) \neq \text{Hom}(X_0, Y_0)$  for some  $X_0, Y_0 \in \text{Ob}(\mathcal{C})$  be given. Consider the sub-category  $B\mathcal{C}$  defined as follows:

$$\text{Ob}(B\mathcal{C}) := \text{Ob}(\mathcal{C}), \text{Hom}_{B\mathcal{C}}(X, Y) := \text{Isom}_{\mathcal{C}}(X, Y).$$

Then this is a sub-category that is **not** full.

